

AAP Guidelines for the Febrile Neonate

References

Aronson PL, Thurm C, Alpern ER, et al; Febrile Young Infant Research Collaborative. Variation in care of the febrile young infant

Blaschke AJ, Korgnski EK, Byington CL. Meningitis in well-appearing febrile infants aged 1–90 days. *Open Forum Infect Dis*. 2018;5(suppl 1):S133

Greenhow TL, Hung YY, Pantell RH. Management and outcomes of previously healthy, full-term, febrile infants ages 7 to 90 days. *Pediatrics*. 2016;138(6):e20160270

Pantell RH, Newman TB, Bernzweig J, et al. Management and outcomes of care of fever in early infancy. *JAMA*. 2004;291(10):1203–1212

Powell EC, Mahajan PV, Roosevelt G, et al; Febrile Infant Working Group of the Pediatric Emergency Care Applied Research Network (PECARN). Epidemiology of bacteremia in febrile infants aged 60 days and younger. *Ann Emerg Med*. 2018;71(2):211–216

Prymula R, Siegrist C-A, Chlibek R, et al. Effect of prophylactic paracetamol administration at time of vaccination on febrile reactions and antibody responses in children: two open-label, randomised controlled trials. *Lancet*. 2009;374(9698):1339–1350