

What is the Future of Surfactant Delivery?

References

- Cummings JJ, Gerday E, Minton S, Katheria A, Albert G, et al; AERO-02 STUDY INVESTIGATORS. Aerosolized Calfactant for Newborns With Respiratory Distress: A Randomized Trial. *Pediatrics*. 2020 Nov;146(5):e20193967. doi: 10.1542/peds.2019-3967. Epub 2020 Oct 15. PMID: 33060258.
- Dargaville PA, Kamlin COF, Orsini F, Wang X, De Paoli AG, et al; OPTIMIST-A Trial Investigators. Effect of Minimally Invasive Surfactant Therapy vs Sham Treatment on Death or Bronchopulmonary Dysplasia in Preterm Infants With Respiratory Distress Syndrome: The OPTIMIST-A Randomized Clinical Trial. *JAMA*. 2021 Dec 28;326(24):2478-2487. doi: 10.1001/jama.2021.21892. PMID: 34902013; PMCID: PMC8715350.
- Gallup JA, Ndakor SM, Pezzano C, Pinheiro JMB. Randomized Trial of Surfactant Therapy via Laryngeal Mask Airway Versus Brief Tracheal Intubation in Neonates Born Preterm. *J Pediatr*. 2023 Mar;254:17-24.e2. doi: 10.1016/j.jpeds.2022.10.009. Epub 2022 Oct 12. PMID: 36241051.
- Gelfand CA, Sakurai R, Wang Y, Liu Y, Segal R, Rehan VK. Inhaled vitamin A is more effective than intramuscular dosing in mitigating hyperoxia-induced lung injury in a neonatal rat model of bronchopulmonary dysplasia. *Am J Physiol Lung Cell Mol Physiol*. 2020 Sep 1;319(3):L576-L584. doi: 10.1152/ajplung.00266.2020. Epub 2020 Aug 5. PMID: 32755324; PMCID: PMC7642870.
- Gupta BK, Saha AK, Mukherjee S, Saha B. Minimally invasive surfactant therapy versus InSurE in preterm neonates of 28 to 34 weeks with respiratory distress syndrome on non-invasive positive pressure ventilation-a randomized controlled trial. *Eur J Pediatr*. 2020 Aug;179(8):1287-1293. doi: 10.1007/s00431-020-03682-9. Epub 2020 May 27. PMID: 32462483; PMCID: PMC7251045.
- Guthrie SO, Fort P, Roberts KD. Surfactant Administration Through Laryngeal or Supraglottic Airways. *Neoreviews*. 2021 Oct;22(10):e673-e688. doi: 10.1542/neo.22-10-e673. PMID: 34599065.
- Härtel C, Herting E, Humberg A, Hanke K, Mehler K, et al; German Neonatal Network. Association of Administration of Surfactant Using Less Invasive Methods With Outcomes in Extremely Preterm Infants Less Than 27 Weeks of Gestation. *JAMA Netw Open*. 2022 Aug 1;5(8):e2225810. doi: 10.1001/jamanetworkopen.2022.25810. Erratum in: *JAMA Netw Open*. 2023 Apr 3;6(4):e2312619. PMID: 35943742; PMCID: PMC9364126.
- Isayama T, Iwami H, McDonald S, Beyene J. Association of Noninvasive Ventilation Strategies With Mortality and Bronchopulmonary Dysplasia Among Preterm Infants: A Systematic Review and Meta-analysis. *JAMA*. 2016 Aug 9;316(6):611-24. doi: 10.1001/jama.2016.10708. Erratum in: *JAMA*. 2016 Sep 13;316(10):1116. PMID: 27532916.
- Kribs A, Roll C, Göpel W, Wieg C, Groneck P, et al; NINSAPP Trial Investigators. Nonintubated Surfactant Application vs Conventional Therapy in Extremely Preterm Infants: A Randomized Clinical Trial. *JAMA Pediatr*. 2015 Aug;169(8):723-30. doi: 10.1001/jamapediatrics.2015.0504. PMID: 26053341.

Minocchieri S, Berry CA, Pillow JJ; CureNeb Study Team. Nebulised surfactant to reduce severity of respiratory distress: a blinded, parallel, randomised controlled trial. Arch Dis Child Fetal Neonatal Ed. 2019 May;104(3):F313-F319. doi: 10.1136/archdischild-2018-315051. Epub 2018 Jul 26. Erratum in: Arch Dis Child Fetal Neonatal Ed. 2020 Mar;105(2):e1. PMID: 30049729; PMCID: PMC6764249.

Moya FR, Mazela J, Shore PM, Simonson SG, Segal R, et al; Preterm Neonate Early Respiratory Management Prospective Observational Study investigators. Prospective observational study of early respiratory management in preterm neonates less than 35 weeks of gestation. BMC Pediatr. 2019 May 11;19(1):147. doi: 10.1186/s12887-019-1518-3. PMID: 31078143; PMCID: PMC6511129.

Pinheiro JM, Santana-Rivas Q, Pezzano C. Randomized trial of laryngeal mask airway versus endotracheal intubation for surfactant delivery. J Perinatol. 2016 Mar;36(3):196-201. doi: 10.1038/jp.2015.177. Epub 2015 Dec 3. PMID: 26633145.

Roberts KD, Brown R, Lampland AL, Leone TA, Rudser KD, et al. Laryngeal Mask Airway for Surfactant Administration in Neonates: A Randomized, Controlled Trial. J Pediatr. 2018 Feb;193:40-46.e1. doi: 10.1016/j.jpeds.2017.09.068. Epub 2017 Nov 22. PMID: 29174079.

Singh AJ, Bronshtein V, Khashu M, Lee K, Potts JE, et al. Vitamin A is systemically bioavailable after intratracheal administration with surfactant in an animal model of newborn respiratory distress. Pediatr Res. 2010 Jun;67(6):619-23. doi: 10.1203/PDR.0b013e3181da8fe8. PMID: 20216104.

Sood BG, Thomas R, Delaney-Black V, Xin Y, Sharma A, Chen X. Aerosolized Beractant in neonatal respiratory distress syndrome: A randomized fixed-dose parallel-arm phase II trial. Pulm Pharmacol Ther. 2021 Feb;66:101986. doi: 10.1016/j.pupt.2020.101986. Epub 2020 Dec 16. PMID: 33338661.

Tribolet S, Hennuy N, Snyers D, Lefèbvre C, Rigo V. Analgosedation before Less-Invasive Surfactant Administration: A Systematic Review. Neonatology. 2022;119(2):137-150. doi: 10.1159/000521553. Epub 2022 Feb 4. PMID: 35124678.

Yeh TF, Chen CM, Wu SY, Husan Z, Li TC, et al. Intratracheal Administration of Budesonide/Surfactant to Prevent Bronchopulmonary Dysplasia. Am J Respir Crit Care Med. 2016 Jan 1;193(1):86-95. doi: 10.1164/rccm.201505-0861OC. PMID: 26351971.