

## ***Seizure Management in the NICU***

### **References**

Ahmad KA, Desai SJ, Bennett MM, Ahmad SF, Ng YT, et al. Changing antiepileptic drug use for seizures in US neonatal intensive care units from 2005 to 2014. *J Perinatol.* 2017 Mar;37(3):296-300. doi: 10.1038/jp.2016.206. Epub 2016 Nov 10. PMID: 27831551.

Duerden EG, Guo T, Dodbiba L, Chakravarty MM, Chau V, et al. Midazolam dose correlates with abnormal hippocampal growth and neurodevelopmental outcome in preterm infants. *Ann Neurol.* 2016 Apr;79(4):548-59. doi: 10.1002/ana.24601. Epub 2016 Feb 15. PMID: 26754148.

Favié LMA, Huitema ADR, van den Broek MPH, Rademaker CMA, de Haan TR, et al; PharmaCool study group\*. Lidocaine as treatment for neonatal seizures: Evaluation of previously developed population pharmacokinetic models and dosing regimen. *Br J Clin Pharmacol.* 2020 Jan;86(1):75-84. doi: 10.1111/bcp.14136. Epub 2020 Jan 3. PMID: 31663153; PMCID: PMC6983510.

Kaminiów K, Pająk M, Pająk R, Paprocka J. Pyridoxine-Dependent Epilepsy and Antiquitin Deficiency Resulting in Neonatal-Onset Refractory Seizures. *Brain Sci.* 2021 Dec 31;12(1):65. doi: 10.3390/brainsci12010065. PMID: 35053812; PMCID: PMC8773593.

Mizrahi EM, Kellaway P. Characterization and classification of neonatal seizures. *Neurology.* 1987 Dec;37(12):1837-44. doi: 10.1212/wnl.37.12.1837. PMID: 3683874.

Murray DM, Boylan GB, Ali I, Ryan CA, Murphy BP, Connolly S. Defining the gap between electrographic seizure burden, clinical expression and staff recognition of neonatal seizures. *Arch Dis Child Fetal Neonatal Ed.* 2008 May;93(3):F187-91. doi: 10.1136/adc.2005.086314. Epub 2007 Jul 11. PMID: 17626147.

Painter MJ, Scher MS, Stein AD, Armatti S, Wang Z, et al. Phenobarbital compared with phenytoin for the treatment of neonatal seizures. *N Engl J Med.* 1999 Aug 12;341(7):485-9. doi: 10.1056/NEJM199908123410704. PMID: 10441604.

Pressler RM, Boylan GB, Marlow N, Blennow M, Chiron C, et al; NEonatal seizure treatment with Medication Off-patent (NEMO) consortium. Bumetanide for the treatment of seizures in newborn babies with hypoxic ischaemic encephalopathy (NEMO): an open-label, dose finding, and feasibility phase 1/2 trial. *Lancet Neurol.* 2015 May;14(5):469-77. doi: 10.1016/S1474-4422(14)70303-5. Epub 2015 Mar 10. PMID: 25765333.

Silverstein FS, Jensen FE. Neonatal seizures. *Ann Neurol.* 2007 Aug;62(2):112-20. doi: 10.1002/ana.21167. PMID: 17683087.

Soul JS, Bergin AM, Stopp C, Hayes B, Singh A, et al; Boston Bumetanide Trial Group. A Pilot Randomized, Controlled, Double-Blind Trial of Bumetanide to Treat Neonatal Seizures. *Ann Neurol.* 2021 Feb;89(2):327-340. doi: 10.1002/ana.25959. Epub 2020 Dec 3. PMID: 33201535; PMCID: PMC8122513.

van den Broek MP, Huitema AD, van Hasselt JG, Groenendaal F, Toet MC, et al. Lidocaine (lignocaine) dosing regimen based upon a population pharmacokinetic model for preterm and term neonates with seizures. *Clin Pharmacokinet*. 2011 Jul;50(7):461-9. doi: 10.2165/11589160-000000000-00000. PMID: 21651313.

van den Broek MP, Rademaker CM, van Straaten HL, Huitema AD, Toet MC, et al. Anticonvulsant treatment of asphyxiated newborns under hypothermia with lidocaine: efficacy, safety and dosing. *Arch Dis Child Fetal Neonatal Ed*. 2013 Jul;98(4):F341-5. doi: 10.1136/archdischild-2012-302678. Epub 2013 Jan 8. PMID: 23303304.

Vasudevan C, Levene M. Epidemiology and aetiology of neonatal seizures. *Semin Fetal Neonatal Med*. 2013 Aug;18(4):185-91. doi: 10.1016/j.siny.2013.05.008. Epub 2013 Jun 5. PMID: 23746578.

Vesoulis ZA, Inder TE, Woodward LJ, Buse B, Vavasseur C, Mathur AM. Early electrographic seizures, brain injury, and neurodevelopmental risk in the very preterm infant. *Pediatr Res*. 2014 Apr;75(4):564-9. doi: 10.1038/pr.2013.245. Epub 2013 Dec 23. PMID: 24366515; PMCID: PMC3961524.