

## ***TeleNeurology in the NICU: Why, How, & What***

### **References**

Bednarek N, Mathur A, Inder T, Wilkinson J, Neil J, Shimony J. Impact of therapeutic hypothermia on MRI diffusion changes in neonatal encephalopathy. *Neurology*. 2012 May 1;78(18):1420-7. doi: 10.1212/WNL.0b013e318253d589. Epub 2012 Apr 18. PMID: 22517107; PMCID: PMC3345786.

Caravale B, Allemand F, Libenson MH. Factors predictive of seizures and neurologic outcome in perinatal depression. *Pediatr Neurol*. 2003 Jul;29(1):18-25. doi: 10.1016/s0887-8994(03)00046-8. PMID: 13679117.

Herman ST, Abend NS, Bleck TP, Chapman KE, Drislane FW, et al; Critical Care Continuous EEG Task Force of the American Clinical Neurophysiology Society. Consensus statement on continuous EEG in critically ill adults and children, part II: personnel, technical specifications, and clinical practice. *J Clin Neurophysiol*. 2015 Apr;32(2):96-108. doi: 10.1097/WNP.000000000000165. PMID: 25626777; PMCID: PMC4434600.

Hunt RW, Liley HG, Wagh D, Schembri R, Lee KJ, et al; Newborn Electrographic Seizure Trial Investigators. Effect of Treatment of Clinical Seizures vs Electrographic Seizures in Full-Term and Near-Term Neonates: A Randomized Clinical Trial. *JAMA Netw Open*. 2021 Dec 1;4(12):e2139604. doi: 10.1001/jamanetworkopen.2021.39604. PMID: 34919132; PMCID: PMC8683963.

Kato T, Okumura A, Hayakawa F, Tsuji T, Hayashi S, et al. Prolonged EEG depression in term and near-term infants with hypoxic ischemic encephalopathy and later development of West syndrome. *Epilepsia*. 2010 Dec;51(12):2392-6. doi: 10.1111/j.1528-1167.2010.02727.x. Epub 2010 Sep 30. PMID: 20887363.

Liu X, Jary S, Cowan F, Thoresen M. Reduced infancy and childhood epilepsy following hypothermia-treated neonatal encephalopathy. *Epilepsia*. 2017 Nov;58(11):1902-1911. doi: 10.1111/epi.13914. Epub 2017 Sep 29. PMID: 28961316.

Lugli L, Balestri E, Berardi A, Guidotti I, Cavalleri F, et al. Brain cooling reduces the risk of postneonatal epilepsy in newborns affected by moderate to severe hypoxic-ischemic encephalopathy. *Minerva Pediatr (Torino)*. 2021 Apr;73(2):150-158. doi: 10.23736/S2724-5276.18.05224-6. Epub 2018 Jul 2. PMID: 29968450.

Philippi H, Wohlrab G, Bettendorf U, Borusiak P, Kluger G, et al. Electroencephalographic evolution of hypsarrhythmia: toward an early treatment option. *Epilepsia*. 2008 Nov;49(11):1859-64. doi: 10.1111/j.1528-1167.2008.01715.x. Epub 2008 Jul 9. PMID: 18631366.

Ruijter BJ, Keijzer HM, Tjepkema-Cloostermans MC, Blans MJ, Beishuizen A, et al; TELSTAR Investigators. Treating Rhythmic and Periodic EEG Patterns in Comatose Survivors of Cardiac Arrest. *N Engl J Med*. 2022 Feb 24;386(8):724-734. doi: 10.1056/NEJMoa2115998. PMID: 35196426.

Rutherford M, Ramenghi LA, Edwards AD, Brocklehurst P, Halliday H, et al. Assessment of brain tissue injury after moderate hypothermia in neonates with hypoxic-ischaemic encephalopathy: a nested substudy of a randomised controlled trial. *Lancet Neurol*. 2010 Jan;9(1):39-45. doi: 10.1016/S1474-4422(09)70295-9. Epub 2009 Nov 5. PMID: 19896902; PMCID: PMC2795146.

Shiple L, Mistry A, Sharkey D. Outcomes of neonatal hypoxic-ischaemic encephalopathy in centres with and without active therapeutic hypothermia: a nationwide propensity score-matched analysis. *Arch Dis Child Fetal Neonatal Ed.* 2022 Jan;107(1):6-12. doi: 10.1136/archdischild-2020-320966. Epub 2021 May 27. PMID: 34045283.

Takeuchi T, Watanabe K. The EEG evolution and neurological prognosis of neonates with perinatal hypoxia [corrected]. *Brain Dev.* 1989;11(2):115-20. doi: 10.1016/s0387-7604(89)80079-8. Erratum in: *Brain Dev* 1989;11(3):203. PMID: 2712234.

Thoresen M, Tooley J, Liu X, Jary S, Fleming P, et al. Time is brain: starting therapeutic hypothermia within three hours after birth improves motor outcome in asphyxiated newborns. *Neonatology.* 2013;104(3):228-33. doi: 10.1159/000353948. Epub 2013 Sep 12. PMID: 24030160.