

Tiny Lungs, Big Impact: Neuroprotective Respiratory Strategies for the Care of Infants Born at 22 to 23 Weeks' Gestation

References

Beauchene MS, Cunningham AM, Stanford AH, Bischoff AR, Dagle JM, et al. Patent ductus arteriosus (PDA) and response to late surfactant treatment in premature infants. *J Perinatol*. 2023 Oct;43(10):1245-1251. doi: 10.1038/s41372-023-01689-2. Epub 2023 May 9. PMID: 37160976.

Berger JN, Elgin TG, Dagle JM, Klein JM, Colaizy TT. Survival and short-term respiratory outcomes of <750 g infants initially intubated with 2.0 mm vs. 2.5 mm endotracheal tubes. *J Perinatol*. 2022 Feb;42(2):202-208. doi: 10.1038/s41372-021-01227-y. Epub 2021 Oct 21. PMID: 34675371; PMCID: PMC8529572.

Carlo WA, McDonald SA, Fanaroff AA, Vohr BR, Stoll BJ, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Neonatal Research Network. Association of antenatal corticosteroids with mortality and neurodevelopmental outcomes among infants born at 22 to 25 weeks' gestation. *JAMA*. 2011 Dec 7;306(21):2348-58. doi: 10.1001/jama.2011.1752. PMID: 22147379; PMCID: PMC3565238.

Chawla S, Natarajan G, Shankaran S, Carper B, Brion LP, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Neonatal Research Network. Markers of Successful Extubation in Extremely Preterm Infants, and Morbidity After Failed Extubation. *J Pediatr*. 2017 Oct;189:113-119.e2. doi: 10.1016/j.jpeds.2017.04.050. Epub 2017 Jun 7. PMID: 28600154; PMCID: PMC5657557.

Chawla S, Wyckoff MH, Rysavy MA, Patel RM, Chowdhury D, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Neonatal Research Network. Association of Antenatal Steroid Exposure at 21 to 22 Weeks of Gestation With Neonatal Survival and Survival Without Morbidities. *JAMA Netw Open*. 2022 Sep 1;5(9):e2233331. doi: 10.1001/jamanetworkopen.2022.33331. PMID: 36156145; PMCID: PMC9513645.

Colaizy TT, Longmuir S, Gertsch K, Abramoff MD, Klein JM. Use of a Supplemental Oxygen Protocol to Suppress Progression of Retinopathy of Prematurity. *Invest Ophthalmol Vis Sci*. 2017 Feb 1;58(2):887-891. doi: 10.1167/iovs.16-20822. PMID: 28159975.

Colaizy TT, Kummert GJ, Kummert CM, Klein JM. Noninvasive Neurally Adjusted Ventilatory Assist in Premature Infants Postextubation. *Am J Perinatol*. 2017 May;34(6):593-598. doi: 10.1055/s-0036-1596053. Epub 2016 Dec 5. PMID: 27919117.

Deshmukh M, Patole S. Antenatal corticosteroids in impending preterm deliveries before 25 weeks' gestation. *Arch Dis Child Fetal Neonatal Ed*. 2018 Mar;103(2):F173-F176. doi: 10.1136/archdischild-2017-313840. Epub 2017 Dec 5. PMID: 29208662.

Ehret DEY, Edwards EM, Greenberg LT, Bernstein IM, Buzas JS, et al. Association of Antenatal Steroid Exposure With Survival Among Infants Receiving Postnatal Life Support at 22 to 25 Weeks' Gestation. *JAMA Netw Open*. 2018 Oct 5;1(6):e183235. doi: 10.1001/jamanetworkopen.2018.3235. PMID: 30646235; PMCID: PMC6324435.

Elgin TG, Stanford AH, Klein JM. First intention high-frequency jet ventilation for periviable infants. *Curr Opin Pediatr*. 2022 Apr 1;34(2):165-169. doi: 10.1097/MOP.0000000000001104. PMID: 35034079.

Elgin TG, Berger JN, Kaluarachchi DC, Dagle JM, Thomas B, et al. Outcomes of infants with birthweights less than 501 g compared to infants weighing 501-750 g at a center utilizing first intention high frequency jet ventilation. *Front Pediatr*. 2024 Sep 9;12:1392079. doi: 10.3389/fped.2024.1392079. PMID: 39315359; PMCID: PMC11416967.

Gaylord MS, Thieme RE, Woodall DL, Quissell BJ. Predicting mortality in low-birth-weight infants with pulmonary interstitial emphysema. *Pediatrics*. 1985 Aug;76(2):219-24. PMID: 4022695.

Gonzalez F, Harris T, Black P, Richardson P. Decreased gas flow through pneumothoraces in neonates receiving high-frequency jet versus conventional ventilation. *J Pediatr*. 1987 Mar;110(3):464-6. doi: 10.1016/s0022-3476(87)80518-8. PMID: 3819950.

Gupta D, Greenberg RG, Natarajan G, Jani S, Sharma A, et al. Respiratory setback associated with extubation failure in extremely preterm infants. *Pediatr Pulmonol*. 2021 Jul;56(7):2081-2086. doi: 10.1002/ppul.25387. Epub 2021 Apr 5. PMID: 33819392.

Higgins RD, Jobe AH, Koso-Thomas M, Bancalari E, Viscardi RM, et al. Bronchopulmonary Dysplasia: Executive Summary of a Workshop. *J Pediatr*. 2018 Jun;197:300-308. doi: 10.1016/j.jpeds.2018.01.043. Epub 2018 Mar 16. PMID: 29551318; PMCID: PMC5970962.

Jensen EA, Dysart K, Gantz MG, McDonald S, Bamat NA, et al. The Diagnosis of Bronchopulmonary Dysplasia in Very Preterm Infants. An Evidence-based Approach. *Am J Respir Crit Care Med*. 2019 Sep 15;200(6):751-759. doi: 10.1164/rccm.201812-2348OC. PMID: 30995069; PMCID: PMC6775872.

Jensen EA, Edwards EM, Greenberg LT, Soll RF, Ehret DEY, Horbar JD. Severity of Bronchopulmonary Dysplasia Among Very Preterm Infants in the United States. *Pediatrics*. 2021 Jul;148(1):e2020030007. doi: 10.1542/peds.2020-030007. Epub 2021 Jun 2. PMID: 34078747; PMCID: PMC8290972.

Juul SE, Wood TR, Comstock BA, Perez K, Gogcu S, et al; PENUT Consortium. Deaths in a Modern Cohort of Extremely Preterm Infants From the Preterm Erythropoietin Neuroprotection Trial. *JAMA Netw Open*. 2022 Feb 1;5(2):e2146404. doi: 10.1001/jamanetworkopen.2021.46404. PMID: 35129596; PMCID: PMC8822378.

Katz LA, Klein JM. Repeat surfactant therapy for postsurfactant slump. *J Perinatol*. 2006 Jul;26(7):414-22. doi: 10.1038/sj.jp.7211533. Epub 2006 May 18. PMID: 16724122.

Keszler M, Donn SM, Bucciarelli RL, Alverson DC, Hart M, et al. Multicenter controlled trial comparing high-frequency jet ventilation and conventional mechanical ventilation in newborn infants with pulmonary interstitial emphysema. *J Pediatr*. 1991 Jul;119(1 Pt 1):85-93. doi: 10.1016/s0022-3476(05)81046-7. PMID: 1906102.

Keszler M, Modanlou HD, Brudno DS, Clark FI, Cohen RS, et al. Multicenter controlled clinical trial of high-frequency jet ventilation in preterm infants with uncomplicated respiratory distress syndrome. *Pediatrics*. 1997 Oct;100(4):593-9. doi: 10.1542/peds.100.4.593. PMID: 9310511.

Kirpalani H, Ratcliffe SJ, Keszler M, Davis PG, Foglia EE, et al; SAIL Site Investigators. Effect of Sustained Inflation vs Intermittent Positive Pressure Ventilation on Bronchopulmonary Dysplasia or Death Among Extremely Preterm Infants: The SAIL Randomized Clinical Trial. *JAMA*. 2019 Mar 26;321(12):1165-1175. doi: 10.1001/jama.2019.1660. PMID: 30912836; PMCID: PMC6439695.

Mori R, Kusuda S, Fujimura M; Neonatal Research Network Japan. Antenatal corticosteroids promote survival of extremely preterm infants born at 22 to 23 weeks of gestation. *J Pediatr*. 2011 Jul;159(1):110-114.e1. doi: 10.1016/j.jpeds.2010.12.039. Epub 2011 Feb 22. PMID: 21334006.

Odom, MW, Ballard, PW. Chapter 15: Developmental and Hormonal Regulation of the Surfactant System. In: McDonald, J., ed. *Lung Growth and Development*. 1st ed. CRC Press; 1997.

Oei JL, Saugstad OD, Lui K, Wright IM, Smyth JP, et al. Targeted Oxygen in the Resuscitation of Preterm Infants, a Randomized Clinical Trial. *Pediatrics*. 2017 Jan;139(1):e20161452. doi: 10.1542/peds.2016-1452. PMID: 28034908.

Oei JL, Kapadia V, Rabi Y, Saugstad OD, Rook D, et al. Neurodevelopmental outcomes of preterm infants after randomisation to initial resuscitation with lower (FiO₂<0.3) or higher (FiO₂>0.6) initial oxygen levels. An individual patient meta-analysis. *Arch Dis Child Fetal Neonatal Ed*. 2022 Jul;107(4):386-392. doi: 10.1136/archdischild-2021-321565. Epub 2021 Nov 1. PMID: 34725105.

Park CK, Isayama T, McDonald SD. Antenatal Corticosteroid Therapy Before 24 Weeks of Gestation: A Systematic Review and Meta-analysis. *Obstet Gynecol*. 2016 Apr;127(4):715-725. doi: 10.1097/AOG.0000000000001355. PMID: 26959200.

Rallis D, Ben-David D, Woo K, Robinson J, Beadles D, et al. Single center experience with first-intention high-frequency jet vs. volume-targeted ventilation in extremely preterm neonates. *Front Pediatr*. 2024 Jan 4;11:1326668. doi: 10.3389/fped.2023.1326668. PMID: 38239592; PMCID: PMC10794594.

Ramanathan R, Sekar KC, Rasmussen M, Bhatia J, Soll RF. Nasal intermittent positive pressure ventilation after surfactant treatment for respiratory distress syndrome in preterm infants <30 weeks' gestation: a randomized, controlled trial. *J Perinatol*. 2012 May;32(5):336-43. doi: 10.1038/jp.2012.1. Epub 2012 Feb 2. Erratum in: *J Perinatol*. 2012 May;32(5):395. PMID: 22301528.

Rysavy M, Nakamura T, Mehler K, Agren J, McNamara P, Backes C; Tiny Baby Collaborative. Use of 2.0-mm endotracheal tubes for periviable infants. *J Perinatol*. 2022 Sep;42(9):1275-1276. doi: 10.1038/s41372-022-01323-7. Epub 2022 Jan 26. PMID: 35082430; PMCID: PMC9314458.

Shalish W, Keszler M, Kovacs L, Chawla S, Latremouille S, et al. Age at First Extubation Attempt and Death or Respiratory Morbidities in Extremely Preterm Infants. *J Pediatr*. 2023 Jan;252:124-130.e3. doi: 10.1016/j.jpeds.2022.08.025. Epub 2022 Aug 24. PMID: 36027982.

Shukla VV, Souder JP, Imbrock G, Hu M, Rahman AKMF, et al. Hospital and Neurodevelopmental Outcomes in Nano-Preterm Infants Receiving Invasive vs Noninvasive Ventilation at Birth. *JAMA Netw Open*. 2022 Aug 1;5(8):e2229105. doi: 10.1001/jamanetworkopen.2022.29105. PMID: 36036932; PMCID: PMC9425146.

Sindelar R, Nakanishi H, Stanford AH, Colaizy TT, Klein JM. Respiratory management for extremely premature infants born at 22 to 23 weeks of gestation in proactive centers in Sweden, Japan, and USA. *Semin Perinatol*. 2022 Feb;46(1):151540. doi: 10.1016/j.semperi.2021.151540. Epub 2021 Nov 10. PMID: 34872750.

Spitzer AR, Butler S, Fox WW. Ventilatory response to combined high frequency jet ventilation and conventional mechanical ventilation for the rescue treatment of severe neonatal lung disease. *Pediatr Pulmonol*. 1989;7(4):244-50. doi: 10.1002/ppul.1950070410. PMID: 2616248.

Supplemental Therapeutic Oxygen for Prethreshold Retinopathy Of Prematurity (STOP-ROP), a randomized, controlled trial. I: primary outcomes. *Pediatrics*. 2000 Feb;105(2):295-310. doi: 10.1542/peds.105.2.295. PMID: 10654946.

Watkins PL, Dagle JM, Bell EF, Colaizy TT. Outcomes at 18 to 22 Months of Corrected Age for Infants Born at 22 to 25 Weeks of Gestation in a Center Practicing Active Management. *J Pediatr*. 2020 Feb;217:52-58.e1. doi: 10.1016/j.jpeds.2019.08.028. Epub 2019 Oct 9. PMID: 31606151.